



**Bibliografia**

1. Capovilla G, Beccaria F, Minicucci F, Sartori S, Vecchi M and Beghi E. Treatment of convulsive status epilepticus in childhood: recommendations of the Italian League Against Epilepsy (LICE). *Epilepsia*, in press.
2. Capovilla G, Vecchi M, Beccaria F, Sartori S, Minicucci F, Vigeveno F. Trattamento dello stato epilettico convulsivo del bambino: raccomandazioni della Lega Italiana Contro l'Epilessia (LICE). 2012; [http://www.lice.it/pdf/sec\\_pediatico.pdf](http://www.lice.it/pdf/sec_pediatico.pdf)
3. Capovilla G, Striano S and Beccaria F. Changes in Panayiotopoulos syndrome over time. *Epilepsia* 2009; 50, suppl 5: 45-8.
4. Dravet C. The core Dravet syndrome phenotype. *Epilepsia* 2011, suppl 2; 3-9.
5. McIntyre J, Robertson S, Norris E et al. Safety and efficacy of buccal midazolam versus rectal diazepam for emergency treatment of seizures in children: a randomised controlled trial. *Lancet* 2005; 366: 205-210.
6. Mpimbaza A, Ndeezi G, Staedke S, Rosenthal PJ, Byarugaba J. Comparison of buccal midazolam with rectal diazepam in the treatment of prolonged seizures in Ugandan children: a randomized clinical trial. *Pediatrics* 2008; 121: e58-64.
7. National Institute for Health and Clinical Excellence (NICE). The epilepsies: the diagnosis and management of the epilepsies in adults and children in primary and secondary care. NHS Evidence, 2012.
8. Scott RC, Besag FM, Neville BG. Buccal midazolam and rectal diazepam for treatment of prolonged seizures in childhood and adolescence: a randomised trial. *Lancet* 1999; 353: 623-626.
9. Specchio N, Trivisano M, Di Ciompo V, Cappelletti S, Masciarelli G, Volkov J, Fusco L and Vigeveno F. Panayiotopoulos syndrome: A clinical, EEG and neuropsychological study of 93 consecutive patients. *Epilepsia* 2011; 51: 2098-2107.

13A03571

